


LPCC: LHeC miniworkshop
CERN, April 18th 2013

LHeC - Low x Kinematics



for the LHeC Study group, <http://cern.ch/lhec>

1. Introduction.

2. eA at the LHeC and comparison to the LHC.

3. Physics case in eA:


- Inclusive measurements and nuclear PDFs.
- Diffraction.
- Final states: dynamics of QCD radiation and hadronization.

4. Summary and outlook.

CDR, arXiv:1206.2913, J. Phys. G 39 (2012) 075001;
arXiv:1211.4831; arXiv:1211.5102




$$R = \frac{f_{i/A}}{A f_{i/p}} \approx \frac{\text{measured}}{\text{expected if no nuclear effects}}$$

- **Lack of data** $\Rightarrow$ models give vastly different results for the nuclear glue at small scales and x : **problem for benchmarking in HIC.**






Yellow Report on Hard Probes, 2004

- Available DGLAP analysis at NLO show large uncertainties at small scales and x .




NLO analysis



nPDFs:


$$R = \frac{f_{i/A}}{A f_{i/p}} \approx \frac{\text{measured}}{\text{expected if no nuclear effects}}$$

- **Lack of data** $\Rightarrow$ models give vastly different results for the nuclear glue at small scales and x : **problem for benchmarking in HIC.**





Yellow Report on
Hard Probes, 2004

- Available DGLAP analysis at NLO show large uncertainties at small scales and x .





Inclusive J/ψ , I202.I383

Small x and saturation:



- **QCD radiation** of partons when **x decreases** leads to a **large number of partons** (gluons), provided each parton **evolves independently** (linearly, $\Delta[xg] \propto xg$).
- This independent evolution **breaks at high densities** (small x or high mass number A): **non-linear effects** ($gg \rightarrow g$, $\Delta[xg] \propto xg - k(xg)^2$).



Small x and saturation:




- **QCD radiation** of partons when **x decreases** leads to a **large number of partons** (gluons), provided each parton **evolves independently** (linearly, $\Delta[xg] \propto xg$).
- This independent evolution **breaks at high densities** (small x or high mass number A): **non-linear effects** ($gg \rightarrow g$, $\Delta[xg] \propto xg - k(xg)^2$).

Status of small-x physics:


- Three pQCD-based alternatives to describe small-x ep and eA data (differences at moderate $Q^2(>\Lambda^2_{\text{QCD}})$ and small x):
 - DGLAP evolution (fixed order perturbation theory).
 - Resummation schemes: BFKL, CCFM, ABF, CCSS.
 - Saturation (CGC, dipole models).
- **Non-linear effects** (unitarity constraints) are density effects: where? $\Rightarrow$ **two-pronged approach at the LHeC: $\downarrow x / \uparrow A$.**



LHeC Relevance for the HI program:




LHeC Relevance for the HI program:



- Nuclear wave function at small x :
nuclear structure functions.

LHeC Relevance for the HI program:




Gluons from saturated nuclei → Glasma? → QGP → Reconfinement

- Nuclear wave function at small x : **nuclear structure functions.**

- Particle production at the very beginning: **which factorisation in eA?**
- How does the system behave as $\sim$ isotropised so fast?: **initial conditions for plasma formation to be studied in eA.**

LHeC Relevance for the HI program:



Gluons from saturated nuclei → Glasma? → QGP → Reconfinement

- Nuclear wave function at small x : **nuclear structure functions.**

- Particle production at the very beginning: **which factorisation in eA?**
- How does the system behave as $\sim$ isotropised so fast?: **initial conditions for plasma formation to be studied in eA.**

- Probing the medium through energetic particles (jet quenching etc.): **modification of QCD radiation and hadronization in the nuclear medium.**

1. Introduction.

2. eA at the LHeC and comparison to the LHC.

3. Physics case in eA:

- Inclusive measurements and nuclear PDFs.
- Diffraction.
- Final states: dynamics of QCD radiation and hadronization.


4. Summary and outlook.

CDR, arXiv:1206.2913, J. Phys. G 39 (2012) 075001;
arXiv:1211.4831; arXiv:1211.5102

Accelerator:

$$\sqrt{s} \approx 0.8 \text{ TeV/nucleon}$$

electron beam	LR ERL	LR
e- energy at IP[GeV]	60	140
luminosity [$10^{32} \text{ cm}^{-2}\text{s}^{-1}$]	10	0.44
polarization [%]	90	90
bunch population [10^9]	2.0	1.6
e- bunch length [mm]	0.3	0.3
bunch interval [ns]	50	50
transv. emit. $\gamma\epsilon_{x,y}$ [mm]	0.05	0.1
rms IP beam size $\sigma_{x,y}$ [μm]	7	7
e- IP beta funct. $\beta^*_{x,y}$ [m]	0.12	0.14
full crossing angle [mrad]	0	0
geometric reduction H_{hg}	0.91	0.94
repetition rate [Hz]	N/A	10
beam pulse length [ms]	N/A	5
ER efficiency	94%	N/A
average current [mA]	6.6	5.4
tot. wall plug power[MW]	100	100




CDR numbers for luminosity, to be considered now as lower bounds.

Luminosity per nucleon




$$L_{eN} = \begin{cases} 9 \times 10^{31} \text{ cm}^{-2}\text{s}^{-1} & \text{(Nominal Pb)} \\ 1.6 \times 10^{32} \text{ cm}^{-2}\text{s}^{-1} & \text{(Ultimate Pb)} \end{cases}$$

$$\text{eD: } L_{eN} = A L_{eA} > \sim 3 \times 10^{31} \text{ cm}^{-2}\text{s}^{-1}$$

LHeC - Low x Kinematics






LHeC - High Q^2 Kinematics






- Small- x demands 1 degree acceptance.
- Higher luminosity would benefit high- x and Q^2 studies.

Kinematics:






- Small- x demands 1 degree acceptance.
- Higher luminosity would benefit high- x and Q^2 studies.

LHC vs. LHeC:



LHC vs. LHeC:



- The LHeC will explore a region overlapping with the LHC:
- in a cleaner experimental setup;
- on firmer theoretical grounds.

- **PbPb:**

- EW bosons.
- VMs in UPCs.
- Ridge.
- ...

- **pPb:**

- Charged particles.
- Ridge.
- Flow.
- Back-to-back correlations, central-forward and forward-forward?
- EW bosons, DY?
- VMs, HF?
- UPCs?
- Jets and interjet activity?
- ...

LHC studies:


● PbPb:

- EW bosons.
- VMs in UPCs.
- Ridge.
- ...


● pPb:

- Charged particles.
- Ridge.
- Flow.
- Back-to-back correlations, central-forward and forward-forward?
- EW bosons, DY?
- VMs, HF?
- UPCs?
- Jets and interjet activity?
- ...

ALICE, 1210.4520



ALICE, 1210.3615




● PbPb:

- EW bosons.
- VMs in UPCs.
- Ridge.
- ...


● pPb:

- Charged particles.
- Ridge.
- Flow.
- Back-to-back correlations, central-forward and forward-forward?
- EW bosons, DY?
- VMs, HF?
- UPCs?
- Jets and interjet activity?
- ...


CMS, I2I0.5482



ALICE, I2I2.2001





ATLAS, I2I2.5198



LHC studies:

- **PbPb:**


- EW bosons.
- VMs in UPCs.
- Ridge.
- ...



- **pPb:**

- Charged particles.
- Ridge.
- Flow.
- Back-to-back correlations, central-forward and forward-forward?
- EW bosons, DY?
- VMs, HF?
- UPCs?
- Jets and interjet activity?
- ...



ATLAS, I303.2084



LHC studies:

- **PbPb:**


- EW bosons.
- VMs in UPCs.
- Ridge.
- ...



- **pPb:**

- Charged particles.
- Ridge.
- Flow.
- Back-to-back correlations, central-forward and forward-forward?
- EW bosons, DY?
- VMs, HF?
- UPCs?
- Jets and interjet activity?
- ...

ATLAS, I303.2084



- The existence of collective effects in pPb is somewhat unexpected.
- Are they really final state? Do they exist also in eA?

1. Introduction.

2. eA at the LHeC and comparison to the LHC.

3. Physics case in eA:

- Inclusive measurements and nuclear PDFs.
- Diffraction.
- Final states: dynamics of QCD radiation and hadronization.




4. Summary and outlook.

CDR, arXiv:1206.2913, J. Phys. G 39 (2012) 075001;
arXiv:1211.4831; arXiv:1211.5102


eA inclusive: comparison

- Good precision can be obtained for $F_{2(c,b)}$ and F_L at small x (Glauberized 3-5 flavor GBW model, NA '02).

Not optimized!



Note the scale!!!




Nuclear PDFs at small x :

- F_2 data substantially reduce the uncertainties in DGLAP analysis; inclusion of charm, beauty (new!); and F_L (new!) also give constraints.


Nuclear PDFs at small x :

- F_2 data substantially reduce the uncertainties in DGLAP analysis; inclusion of charm, beauty (new!); and F_L (new!) also give constraints.




Nuclear PDFs at small x :



- F_2 data substantially reduce the uncertainties in DGLAP analysis; inclusion of charm, beauty (new!); and F_L (new!) also give constraints.




Diffraction in ep and shadowing:





- Diffraction is linked to nuclear shadowing through basic QFT (Gribov): eD to test and set the 'benchmark' for new effects.




Diffraction in ep and shadowing:





- Diffraction is linked to nuclear shadowing through basic QFT (Gribov): eD to test and set the 'benchmark' for new effects.



LHeC Elastic VM production in eA:





- For the **coherent case**, predictions available.
- **Challenging** experimental problem (neutron tagging in ZDC?).




Transverse scan: elastic VM

- t-differential measurements give a gluon transverse mapping of the hadron/nucleus.




Transverse scan: elastic VM

- t-differential measurements give a gluon transverse mapping of the hadron/nucleus.




- Large extent in t with good precision.
- **Sizable saturation effects expected.**




LHeC Dihadron azimuthal decorrelation:


- Dihadron **azimuthal decorrelation**: currently discussed at RHIC as suggestive of saturation.



- At the LHeC it could be studied far from the kinematical limits.




$$C(\phi_{12}) = \frac{1}{\frac{d\sigma(\gamma^* N \rightarrow h_1 X)}{dz_{h1}}} \frac{d\sigma \gamma^* N \rightarrow h_1 h_2 + X}{dz_{h1} dz_{h2} d\phi_{12}}$$




Radiation and hadronization:

- **LHeC: dynamics of QCD radiation and hadronization.**
- Most relevant for particle production off nuclei and for QGP analysis in HIC.


- **Low energy:** hadronization inside $\rightarrow$ formation time, (pre-)hadronic absorption,...



- **High energy:** partonic evolution altered in the nuclear medium.




$$R_A^h(z, \nu) = \frac{1}{N_A^e} \frac{dN_A^h(z, \nu)}{d\nu dz} \bigg/ \frac{1}{N_D^e} \frac{dN_D^h(z, \nu)}{d\nu dz}$$




Radiation and hadronization:

- **LHeC: dynamics of QCD radiation and hadronization.**
- Most relevant for particle production off nuclei and for QGP analysis in HIC.


- **Low energy:** hadronization inside $\rightarrow$ formation time, (pre-)hadronic absorption,...




- **High energy:** partonic evolution altered in the nuclear medium.



$$R_A^h(z, \nu) = \frac{1}{N_A^e} \frac{dN_A^h(z, \nu)}{d\nu dz} \bigg/ \frac{1}{N_D^e} \frac{dN_D^h(z, \nu)}{d\nu dz}$$



Jets:





- Jets: large E_T even in eA.
- Useful for studies of parton dynamics in nuclei (hard probes), and for photon structure.
- Background subtraction, detailed reconstruction pending.

Summary:

- **At an LHeC@CERN:**

- High-precision tests of collinear factorization(s) and determination of PDFs.
- Unprecedented access to small x in p and A .
- Novel sensitivity to physics beyond standard pQCD.
- Stringent tests of QCD radiation and hadronization.
- Transverse scan of the hadron/nucleus at small x .
- ... with implications on our understanding of QGP.

- **The LHeC will answer the question of saturation/non-linear dynamics. For that, ep AND eA essential!!!**






- With CERN and NuPECC mandate to further motivate the physics case and produce a TDR around 2015, several items have to be done/improved:
 - Refine DGLAP fits with flavour decomposition (include neutrino data, relax assumptions) and optimized F_L scenarios, and LHC data.
 - Monte Carlo generators!!!
 - Studies on diffraction: separation of coherent from incoherent, ndPDFs, dijets,...
 - Large x, EW bosons.
 - Nuclear GPDs: nuclear DVCS etc.
 - eD.
 - Jet reconstruction, angular decorrelation...
 - ...
- Cooperation with EIC in some of these items desirable.




- With CERN and NuPECC mandate to further motivate the physics case and produce a TDR around 2015, several items have to be done/improved:
 - Refine DGLAP fits with flavour decomposition (include neutrino data, relax assumptions) and optimized F_L scenarios, and LHC data.
 - Monte Carlo generators!!!
 - Studies on diffraction: separation of coherent from incoherent, ndPDFs, dijets,...
 - Large x, EW bosons.
 - Nuclear GPDs: nuclear DVCS etc.
 - eD.
 - Jet reconstruction, angular decorrelation...
 - ...
- Thanks for your attention!***
- Cooperation with EIC in some of these items desirable.

Backup:

LHC vs. LHeC:



LHC vs. LHeC:



- The LHeC will explore a region overlapping with the LHC:
- in a cleaner experimental setup;
- on firmer theoretical grounds.

- **PbPb:**

- EW bosons.
- VMs in UPCs.
- Ridge.
- ...

- **pPb:**

- Charged particles.
- Ridge.
- Flow.
- Back-to-back correlations, central-forward and forward-forward?
- EW bosons, DY?
- VMs, HF?
- UPCs?
- Jets and interjet activity?
- ...





LHC studies:

● PbPb:

- EW bosons.
- VMs in UPCs.
- Ridge.
- ...

● pPb:

- Charged particles.
- Ridge.
- Flow.
- Back-to-back correlations, central-forward and forward-forward?
- EW bosons, DY?
- VMs, HF?
- UPCs?
- Jets and interjet activity?
- ...




LHC studies:

- **PbPb:**

- EW bosons.
- VMs in UPCs.
- Ridge.
- ...

- **pPb:**

- Charged particles.
- Ridge.
- Flow.
- Back-to-back correlations, central-forward and forward-forward?
- EW bosons, DY?
- VMs, HF?
- UPCs?
- Jets and interjet activity?
- ...



CMS, I I 05.2438


● PbPb:

- EW bosons.
- VMs in UPCs.
- Ridge.
- ...


● pPb:

- Charged particles.
- Ridge.
- Flow.
- Back-to-back correlations, central-forward and forward-forward?
- EW bosons, DY?
- VMs, HF?
- UPCs?
- Jets and interjet activity?
- ...

(a) CMS $\int L dt = 3.1 \mu\text{b}^{-1}$
PbPb $\sqrt{s_{NN}} = 2.76 \text{ TeV}$, 0-5% centrality




(b) CMS Simulations
PYTHIA8 pp MC $\sqrt{s} = 2.76 \text{ TeV}$




ALICE, I I 09.2501

$3 < p_T^t < 4 \text{ GeV/c}$
 $2 < p_T^a < 2.5 \text{ GeV/c}$
Pb-Pb 2.76 TeV
0-10%



$8 < p_T^t < 15 \text{ GeV/c}$
 $6 < p_T^a < 8 \text{ GeV/c}$
Pb-Pb 2.76 TeV
0-20%



LHC studies:


● PbPb:

- EW bosons.
- VMs in UPCs.
- Ridge.
- ...


● pPb:

- Charged particles.
- Ridge.
- Flow.
- Back-to-back correlations, central-forward and forward-forward?
- EW bosons, DY?
- VMs, HF?
- UPCs?
- Jets and interjet activity?
- ...

ALICE, 1210.4520



ALICE, 1210.3615




● PbPb:

- EW bosons.
- VMs in UPCs.
- Ridge.
- ...


● pPb:

- Charged particles.
- Ridge.
- Flow.
- Back-to-back correlations, central-forward and forward-forward?
- EW bosons, DY?
- VMs, HF?
- UPCs?
- Jets and interjet activity?
- ...

CMS pPb $\sqrt{s_{NN}} = 5.02$ TeV, $N_{trk}^{offline} < 35$
 $1 < p_T < 3$ GeV/c




(a) CMS pPb $\sqrt{s_{NN}} = 5.02$ TeV, $N_{trk}^{offline} \geq 110$
 $1 < p_T < 3$ GeV/c




CMS, I2I0.5482

$2 < p_{T,trig} < 4$ GeV/c
 $1 < p_{T,assoc} < 2$ GeV/c




$2 < p_{T,trig} < 4$ GeV/c
 $1 < p_{T,assoc} < 2$ GeV/c




ALICE, I2I2.2001

ATLAS pPb $\sqrt{s_{NN}} = 5.02$ TeV
 $\Sigma E_T^{Pb} < 20$ GeV
 $\int L \approx 1 \mu b^{-1}$ $0.5 < p_T^{ab} < 4$ GeV



(a)

$\Sigma E_T^{Pb} > 80$ GeV



ATLAS, I2I2.5198

LHC studies:


- **PbPb:**

- EW bosons.
- VMs in UPCs.
- Ridge.
- ...

- **pPb:**

- Charged particles.
- Ridge.
- Flow.
- Back-to-back correlations, central-forward and forward-forward?
- EW bosons, DY?
- VMs, HF?
- UPCs?
- Jets and interjet activity?
- ...

ATLAS, I303.2084




- **PbPb:**

- EW bosons.
- VMs in UPCs.
- Ridge.
- ...

- **pPb:**

- Charged particles.
- Ridge.
- Flow.
- Back-to-back correlations, central-forward and forward-forward?
- EW bosons, DY?
- VMs, HF?
- UPCs?
- Jets and interjet activity?
- ...

ATLAS, I303.2084



- The existence of collective effects in pPb is somewhat unexpected.
- Are they really final state? Do they exist also in eA?

LHeC scenarios:

config.	E(e)	E(N)	N	$\int L(e^+)$	$\int L(e^-)$	Pol	L/ 10^{32}	P/MW	years	type
For F_2										
A	20	7	p	1	1	-	1	10	1	SPL
B	50	7	p	50	50	0.4	25	30	2	RR hiQ ²
C	50	7	p	1	1	0.4	1	30	1	RR lo x
D	100	7	p	5	10	0.9	2.5	40	2	LR
E	150	7	p	3	6	0.9	1.8	40	2	LR
F	50	3.5	D	1	1	--	0.5	30	1	eD
G	50	2.7	Pb	10^{-3}	10^{-3}	0.4	10^{-3}	30	1	ePb
H	50	1	p	--	1	--	25	30	1	lowEp
I	50	3.5	Ca	$5 \cdot 10^{-3}$		?	$5 \cdot 10^{-3}$	?	?	eCa


- For F_L : 10, 25, 50 + 2750 (7000); $Q^2 \leq s_x$; Lumi=5, 10, 100 pb⁻¹ respectively; charm and beauty: same efficiencies in ep and eA.

Note: F_L in eA


$$\sigma_r^{NC} = \frac{Q^4 x}{2\pi\alpha^2 Y_+} \frac{d^2\sigma^{NC}}{dx dQ^2} = F_2 \left[1 - \frac{y^2}{Y_+} \frac{F_L}{F_2} \right], \quad Y_+ = 1 + (1 - y)^2$$

- F_L traces the nuclear effects on the glue (Cazarotto et al '08).
- Uncertainties in the extraction of F_2 due to the unknown nuclear effects on F_L of order 5 % (larger than expected stat.+syst.) $\Rightarrow$ measure F_L or use the reduced cross section (but then ratios at two energies...).




NA, Paukkunen, Salgado, Tywoniuk, '10





Elastic VM production in ep:




- Elastic J/ψ production appears as a candidate to signal saturation effects at work!!!






- **Challenging** experimental problem, requires Monte Carlo simulation with detailed understanding of the nuclear break-up.
- For the **coherent case**, predictions available.




Radiation and hadronization:

- Large (NLO) yields at small- x (H1 cuts, 3 times higher if relaxed).
- Nuclear effects in hadronization at small ν (LO plus QW, Arleo '03).





$$R_A^h(z, \nu) = \frac{1}{N_A^e} \frac{dN_A^h(z, \nu)}{d\nu dz} \bigg/ \frac{1}{N_D^e} \frac{dN_D^h(z, \nu)}{d\nu dz}$$




Radiation and hadronization:

- Large (NLO) yields at small- x (HI cuts, 3 times higher if relaxed).
- Nuclear effects in hadronization at small v (LO plus QW, Arleo '03).






Radiation and hadronization:

- Large (NLO) yields at small- x (HI cuts, 3 times higher if relaxed).
- Nuclear effects in hadronization at small ν (LO plus QW, Arleo '03).



$$\frac{1}{N_D^e} \frac{dN_D^h(z, \nu)}{d\nu dz}$$

$$R_A^h(z, \nu) = \frac{1}{N_A^e} \frac{dN_A^h(z, \nu)}{d\nu dz}$$



- With CERN and NuPECC mandate to further motivate the physics case and produce a TDR around 2015, several items have to be done/improved:
 - Refine DGLAP fits with flavour decomposition (include neutrino data, relax assumptions) and optimized F_L scenarios, and LHC data.
 - Monte Carlo generators!!!
 - Studies on diffraction: separation of coherent from incoherent, ndPDFs, dijets,...
 - Large x, EW bosons.
 - Nuclear GPDs: nuclear DVCS etc.
 - eD.
 - Jet reconstruction, angular decorrelation...
 - ...
- Cooperation with EIC in some of these items desirable.

- With CERN and NuPECC make physics case and produce a TDR to be done/improved:

- Refine DGLAP fits with flavour data, relax assumptions) and optimise Monte Carlo generators!!!
- Studies on diffraction: separate PDFs, dijets,...
- Large x, EW bosons.
- Nuclear GPDs: nuclear DVCS
- eD.
- Jet reconstruction, angular deconvolution
- ...

→ Cooperation with EIC in some areas

2. Recommendations and Roadmap


NuPECC LRP

EURISOL in future updates of the ESFRI list, based on the successful EURISOL Design Study in FP6.

- The Technical Design Study for intense radioactive ion beams at ISOL@MYRRHA.
- The Technical Design Study for a polarised proton-antiproton, PAX, and an electron-nucleon/ion collider, ENC, at FAIR.
- The Technical Design Study for a high-energy electron-proton/ion collider, LHeC, at CERN.
- The inclusion of Nuclear Physics programmes at the multi-purpose facilities ELI and ESS.

2.2 Facilities Roadmap

We present below the roadmap for building new large-scale Nuclear Physics research infrastructures in Europe. The time span ranges until the middle of the next decade. Facilities whose first phases have already been approved are coloured in blue, future upgrades thereof in dark blue. The ISOL facilities SPIRAL 2, HIE-ISOLDE and SPES are designated to lead to EURISOL. PAX and the ENC at FAIR, EURISOL and the LHeC at CERN are still in the design or R&D phase. They are coloured in purple.



Roadmap for New Large Scale Facilities.

- With CERN and NuPECC mandate to further motivate the physics case and produce a TDR around 2015, several items have to be done/improved:
 - Refine DGLAP fits with flavour decomposition (include neutrino data, relax assumptions) and optimized F_L scenarios, and LHC data.
 - Monte Carlo generators!!!
 - Studies on diffraction: separation of coherent from incoherent, ndPDFs, dijets,...
 - Large x, EW bosons.
 - Nuclear GPDs: nuclear DVCS etc.
 - eD.
 - Jet reconstruction, angular decorrelation...
 - ...
- Cooperation with EIC in some of these items desirable.

- With CERN and NuPECC mandate to further motivate the physics case and produce a TDR around 2015, several items have to be done/improved:
 - Refine DGLAP fits with flavour decomposition (include neutrino data, relax assumptions) and optimized F_L scenarios, and LHC data.
 - Monte Carlo generators!!!
 - Studies on diffraction: separation of coherent from incoherent, ndPDFs, dijets,...
 - Large x, EW bosons.
 - Nuclear GPDs: nuclear DVCS etc.
 - eD.
 - Jet reconstruction, angular decorrelation...
 - ...
- Thanks for your attention!***
- Cooperation with EIC in some of these items desirable.