

Outline

■ Relativistic Kinematics

- ▶ (4-momentum)² invariance, invariant mass
- ▶ Hypothesis testing, production thresholds
- ▶ Cross-sections, flux and luminosity, accelerators
- ▶ Particle lifetime, decay length, width

Today

■ Classification of particles

- ▶ Fermions and bosons
- ▶ Leptons, hadrons, quarks
- ▶ Mesons, baryons

■ Quark Model

- ▶ Meson and baryon multiplets
- ▶ Isospin, strangeness, c, b, t quarks

■ Particle Interactions

- ▶ Colour charge, QCD, gluons, fragmentation, **running couplings**
- ▶ **Strong and weak decays, conservation rules**
- ▶ Virtual particles and range of forces
- ▶ Parity, charge conjugation, CP
- ▶ Weak decays of quarks
- ▶ Charmonium and upsilon systems

Previous
lecture

■ Electroweak Interactions

- ▶ Charged and neutral currents
- ▶ W, Z, LEP experiments
- ▶ Higgs and the future



■ LHC Experiments

■ Future - introduction to accelerator physics

- [Lecture 11 \(4 slides/page\)](#) - Fragmentation, running couplings, particle decays
 - Halzen and Martin, pp. 16-26
 - Griffiths, pp. 298-301
 - Williams, pp. 221-227
 - Perkins, pp. 44-46

α_s Summary


Consistent value of α_s measured in many different reactions. Note that values are all transformed (“evolved”) to a **single energy scale** to allow comparison, using “Renormalisation Group Equations”. QCD predicts how α_s varies with energy, not its actual value




[K. Nakamura *et al.*](#) (Particle Data Group), J. Phys. G 37, 075021 (2010)
[\[http://pdg.lbl.gov/2011/reviews/rpp2011-rev-qcd.pdf\]](http://pdg.lbl.gov/2011/reviews/rpp2011-rev-qcd.pdf)

Running Coupling Constants

- Coupling “constants” are said to “run” (change their strength) with energy
- For **electromagnetism**, the coupling “constant”, α_{EM} , **increased** with energy
- For **weak force** the coupling constant **decreases** with energy
 - ▶ E.M. and weak merge at ~ 100 GeV: “electroweak unification”
- For **strong force** coupling, α_s , **decreases** with energy




Requires gauge coupling unification
Low energy SUSY
Nucleon decay



[Allanach, Blair, Kraml, Martyn, Polesello,
Porod, Zerwas, LHC-LC Report, p339]

Strong, e.m., weak interactions (W.I.)

- So far, have discussed strong interaction in terms of binding quarks into hadrons
- Particle decays also determined by type of interactions allowed
- **Strength** of interaction reflected in **lifetime** of decaying particle
- Many hadronic resonances, lifetimes
 - ▶ $\tau \sim 10^{-23}\text{s}$
 - ▶ Deduced from width, $\Gamma \sim 10\text{-}100\text{ MeV}$
 - ▶ These are **Strong Interaction** decays
- Some much longer lived hadrons
 - ▶ $\tau \sim 10^{-10}\text{s}$
 - ▶ Can be measured directly
 - ▶ These are **Weak Interaction** decays
- Some with intermediate lifetimes (e.m.)



Conservation Rules

| Interaction | Symbol | SI | EM | WI |
|----------------------------|--------|----|----|----|
| Energy | E | ✓ | ✓ | ✓ |
| Momentum | P | ✓ | ✓ | ✓ |
| Angular Mom ⁿ . | J | ✓ | ✓ | ✓ |
| Charge (e.m, colour) | Q | ✓ | ✓ | ✓ |
| Fermion number | | ✓ | ✓ | ✓ |
| Quark number | | ✓ | ✓ | ✓ |
| Baryon number | B | ✓ | ✓ | ✓ |
| Lepton number | L | ✓ | ✓ | ✓ |
| Electron number | L_e | ✓ | ✓ | ✓ |
| Muon number | L_m | ✓ | ✓ | ✓ |
| Tau number | L_t | ✓ | ✓ | ✓ |
| Quark flavour | | ✓ | ✓ | ✗ |
| Isospin | I | ✓ | ✗ | ✗ |
| Parity | P | ✓ | ✓ | ✗ |
| Charge Conjugation | C | ✓ | ✓ | ✗ |
| Time reversal | T | ✓ | ✓ | ✗ |
| Matter-Antimatter | CP | ✓ | ✓ | ✗ |
| Quantum Field Theory | CPT | ✓ | ✓ | ✓ |

✓ conserved
 ✗ Not necessarily conserved

[For info.] Running Couplings

EM case

$$\alpha_{EM}(|q^2|) = \frac{\alpha(0)}{1 - \left(\frac{\alpha(0)}{3\pi}\right) \ln(|q^2|/m^2)} \quad |q^2| \gg m^2$$

QCD case

$$\alpha_S(|q^2|) = \frac{\alpha_S(\mu^2)}{1 + \left(\frac{\alpha_S(\mu^2)}{12\pi}\right) [11N_{colours} - 2N_{flavours}] \ln(|q^2|/m^2)} \quad |q^2| \gg |\mu^2|$$