

Outline

■ Relativistic Kinematics

- ▶ (4-momentum)² invariance, invariant mass
- ▶ Hypothesis testing, production thresholds
- ▶ Cross-sections, flux and luminosity, accelerators
- ▶ Particle lifetime, decay length, width

■ Classification of particles

- ▶ Fermions and bosons
- ▶ Leptons, hadrons, quarks
- ▶ Mesons, baryons

■ Quark Model

- ▶ Meson and baryon multiplets
- ▶ Isospin, strangeness, c, b, t quarks

■ Particle Interactions

- ▶ Colour charge, QCD, gluons, fragmentation, **running couplings**
- ▶ **Strong and weak decays, conservation rules**
- ▶ Virtual particles and range of forces
- ▶ Parity, charge conjugation, CP
- ▶ Weak decays of quarks
- ▶ Charmonium and upsilon systems

■ Electroweak Interactions

- ▶ Charged and neutral currents
- ▶ W, Z, LEP experiments
- ▶ Higgs and the future

■ LHC Experiments

■ Future - introduction to accelerator physics

Today



- [Lecture 12 \(4 slides/page\)](#)
 - Griffiths, pp. 66-71, 79-82, 84-85.

Previous
lecture

- [Lecture 11 \(4 slides/page\)](#) - Fragmentation, running couplings, particle decays
 - Halzen and Martin, pp. 16-26
 - Griffiths, pp. 298-301
 - Williams, pp. 221-227
 - Perkins, pp. 44-46

α_s Summary

Consistent value of α_s measured in many different reactions. Note that values are all transformed (“evolved”) to a **single energy scale** to allow comparison, using “Renormalisation Group Equations”. QCD predicts how α_s varies with energy, not its actual value



“Confinement”

“asymptotic freedom”


Long range (low energy)

Short range (high energy)


[K. Nakamura *et al.* \(Particle Data Group\), J. Phys. G 37, 075021 \(2010\)](http://pdg.lbl.gov/2011/reviews/rpp2011-rev-qcd.pdf)
[\[http://pdg.lbl.gov/2011/reviews/rpp2011-rev-qcd.pdf\]](http://pdg.lbl.gov/2011/reviews/rpp2011-rev-qcd.pdf)

Running Coupling Constants

- Coupling “constants” are said to “run” (change their strength) with energy
- For **electromagnetism**, the coupling “constant”, α_{EM} , **increased** with energy
- For **weak force** the coupling constant **decreases** with energy
 - ▶ E.M. and weak merge at ~ 100 GeV: “electroweak unification”
- For **strong force** coupling, α_s , **decreases** with energy




Requires gauge coupling unification
Low energy SUSY
Nucleon decay



[Allanach, Blair, Kraml, Martyn, Polesello,
Porod, Zerwas, LHC-LC Report, p339]

Strong, e.m., weak interactions (W.I.)

- So far, have discussed strong interaction in terms of binding quarks into hadrons
- Particle decays also determined by type of interactions allowed
- **Strength** of interaction reflected in **lifetime** of decaying particle
- Many hadronic resonances, lifetimes
 - ▶ $\tau \sim 10^{-23} \text{s}$
 - ▶ **Deduced from width,**
 $\Gamma \sim 10\text{-}100 \text{ MeV}$
 - ▶ **These are Strong Interaction decays**
- Some much longer lived hadrons
 - ▶ $\tau \sim 10^{-10} \text{s}$
 - ▶ **Can be measured directly**
 - ▶ **These are Weak Interaction decays**
- Some with intermediate lifetimes (e.m.)



Conservation Rules

Interaction	Symbol	SI	EM	WI
Energy	E	✓	✓	✓
Momentum	P	✓	✓	✓
Angular Mom ⁿ .	J	✓	✓	✓
Charge (e.m, colour)	Q	✓	✓	✓
Fermion number		✓	✓	✓
Quark number		✓	✓	✓
Baryon number	B	✓	✓	✓
Lepton number	L	✓	✓	✓
Electron number	L_e	✓	✓	✓
Muon number	L_m	✓	✓	✓
Tau number	L_t	✓	✓	✓
Quark flavour		✓	✓	✗
Isospin	I	✓	✗	✗
Parity	P	✓	✓	✗
Charge Conjugation	C	✓	✓	✗
Time reversal	T	✓	✓	✗
Matter-Antimatter	CP	✓	✓	✗
Quantum Field Theory	CPT	✓	✓	✓

✓	conserved
	Not
✗	necessarily
	conserved

[For info.] Running Couplings

EM case

$$\alpha_{EM}(|q^2|) = \frac{\alpha(0)}{1 - \left(\frac{\alpha(0)}{3\pi}\right) \ln(|q^2|/m^2)} \quad |q^2| \gg m^2$$

QCD case

$$\alpha_S(|q^2|) = \frac{\alpha_S(\mu^2)}{1 + \left(\frac{\alpha_S(\mu^2)}{12\pi}\right) [11N_{colours} - 2N_{flavours}] \ln(|q^2|/m^2)} \quad |q^2| \gg |\mu^2|$$