

Outline

- Relativistic Kinematics
 - ▶ (4-momentum)² invariance, invariant mass
 - ▶ Hypothesis testing, production thresholds
 - ▶ Cross-sections, flux and luminosity, accelerators
 - ▶ Particle lifetime, decay length, width
- Classification of particles
 - ▶ Fermions and bosons
 - ▶ Leptons, hadrons, quarks
 - ▶ Mesons, baryons
- Quark Model
 - ▶ Meson and baryon multiplets
 - ▶ Isospin, strangeness, c, b, t quarks
- Particle Interactions
 - ▶ Colour charge, QCD, gluons, fragmentation, running couplings
 - ▶ Strong and weak decays, conservation rules
 - ▶ Virtual particles and range of forces
 - ▶ Parity, charge conjugation, CP
 - ▶ Weak decays of quarks
 - ▶ Charmonium and upsilon systems
- Electroweak Interactions
 - ▶ Charged and neutral currents
 - ▶ W, Z, LEP experiments
 - ▶ Higgs and the future
- LHC Experiments
- Future - introduction to accelerator physics



Today

Previous lecture

- Lecture 11 (4 slides/page) - Fragmentation, running couplings, particle decays
 - Halzen and Martin, pp. 16-26
 - Griffiths, pp. 298-301
 - Williams, pp. 221-227
 - Perkins, pp. 44-46

α_s Summary


Consistent value of α_s measured in many different reactions. Note that values are all transformed ("evolved") to a single energy scale to allow comparison, using "Renormalisation Group Equations". QCD predicts how α_s varies with energy, not its actual value




K. Nakamura et al. (Particle Data Group), J. Phys. G 37, 075021 (2010)
[\[http://pdg.lbl.gov/2011/reviews/rpp2011-rev-qcd.pdf\]](http://pdg.lbl.gov/2011/reviews/rpp2011-rev-qcd.pdf)

Running Coupling Constants

- Coupling "constants" are said to "run" (change their strength) with energy
- For **electromagnetism**, the coupling "constant", α_{EM} , **increased** with energy
- For **weak force** the coupling constant **decreases** with energy
 - ▶ E.M. and weak merge at ~ 100 GeV: "electroweak unification"
- For **strong force** coupling, α_s , **decreases** with energy




Requires gauge coupling unification
 Low energy SUSY
 Nucleon decay



[Allanach, Blair, Kraml, Martyn, Polesello, Porod, Zerwas, LHC-LC Report, p339]

Strong, e.m., weak interactions (W.I.)

- So far, have discussed strong interaction in terms of binding quarks into hadrons
- Particle decays also determined by type of interactions allowed
- Strength** of interaction reflected in **lifetime** of decaying particle
- Many hadronic resonances, lifetimes
 - $\tau \sim 10^{-23}$ s
 - Deduced from width, $\Gamma \sim 10-100$ MeV
 - These are **Strong Interaction** decays
- Some **much longer lived** hadrons
 - $\tau \sim 10^{-10}$ s
 - Can be measured directly
 - These are **Weak Interaction** decays
- Some with **intermediate** lifetimes (e.m.)



Conservation Rules

Interaction	Symbol	SI	EM	WI
Energy	E	✓	✓	✓
Momentum	P	✓	✓	✓
Angular Mom ⁿ	J	✓	✓	✓
Charge (e.m, colour)	Q	✓	✓	✓
Fermion number		✓	✓	✓
Quark number		✓	✓	✓
Baryon number	B	✓	✓	✓
Lepton number	L	✓	✓	✓
Electron number	L _e	✓	✓	✓
Muon number	L _μ	✓	✓	✓
Tau number	L _τ	✓	✓	✓
Quark flavour		✓	✓	✗
Isospin	I	✓	✗	✗
Parity	P	✓	✓	✗
Charge Conjugation	C	✓	✓	✗
Time reversal	T	✓	✓	✗
Matter-Antimatter	CP	✓	✓	✗
Quantum Field Theory	CPT	✓	✓	✓

✓ conserved
✗ Not necessarily conserved

[For info.] Running Couplings

EM case

$$\alpha_{EM}(q^2) = \frac{\alpha(0)}{1 - \left(\frac{\alpha(0)}{3\pi}\right) \ln(q^2/m^2)} \quad |q^2| \gg m^2$$

QCD case

$$\alpha_S(q^2) = \frac{\alpha_S(\mu^2)}{1 + \left(\frac{\alpha_S(\mu^2)}{12\pi}\right) [11N_{colours} - 2N_{flavours}] \ln(q^2/m^2)} \quad |q^2| \gg |\mu^2|$$