

# Outline

- Relativistic Kinematics
  - ▶ (4-momentum)<sup>2</sup> invariance, invariant mass
  - ▶ Hypothesis testing, production thresholds
  - ▶ Cross-sections, flux and luminosity, accelerators
  - ▶ Particle lifetime, decay length, width
- Classification of particles
  - ▶ Fermions and bosons
  - ▶ Leptons, hadrons, quarks
  - ▶ Mesons, baryons
- Quark Model
  - ▶ Meson and baryon multiplets
  - ▶ Isospin, strangeness, c, b, t quarks
- Particle Interactions
  - ▶ Colour charge, QCD, gluons, fragmentation, running couplings
  - ▶ Strong and weak decays, conservation rules
  - ▶ Virtual particles and range of forces
  - ▶ Parity, charge conjugation, CP
  - ▶ Weak decays of quarks
  - ▶ Charmonium and upsilon systems
- Electroweak Interactions
  - ▶ Charged and neutral currents
  - ▶ W, Z, LEP experiments
  - ▶ Higgs and the future
- LHC Experiments
- Future - introduction to accelerator physics

Today



• Lecture 12 (4 slides/page)  
o Griffiths, pp. 66-71, 79-82, 84-85.

Previous lecture

• Lecture 11 (4 slides/page) - Fragmentation, running couplings, particle decays  
o Halzen and Martin, pp. 16-26  
o Griffiths, pp. 298-301  
o Williams, pp. 221-227  
o Perkins, pp. 44-46

# $\alpha_s$ Summary


Consistent value of  $\alpha_s$  measured in many different reactions. Note that values are all transformed ("evolved") to a single energy scale to allow comparison, using "Renormalisation Group Equations". QCD predicts how  $\alpha_s$  varies with energy, not its actual value




K. Nakamura et al. (Particle Data Group), J. Phys. G 37, 075021 (2010)  
[http://pdg.lbl.gov/2011/reviews/rpp2011-rev-qcd.pdf]

# Running Coupling Constants

- Coupling "constants" are said to "run" (change their strength) with energy
- For **electromagnetism**, the coupling "constant",  $\alpha_{EM}$ , **increased** with energy
- For **weak force** the coupling constant **decreases** with energy
  - ▶ E.M. and weak merge at  $\sim 100$  GeV: "electroweak unification"
- For **strong force** coupling,  $\alpha_s$ , **decreases** with energy




Requires gauge coupling unification  
Low energy SUSY  
Nucleon decay



[Allanach, Blair, Kraml, Martyn, Polesello, Porod, Zerwas, LHC-LC Report, p339]

## Strong, e.m., weak interactions (W.I.)

- So far, have discussed strong interaction in terms of binding quarks into hadrons
- Particle decays also determined by type of interactions allowed
- Strength** of interaction reflected in **lifetime** of decaying particle
- Many hadronic resonances, lifetimes
  - $\tau \sim 10^{-23}$ s
  - Deduced from width,  $\Gamma \sim 10-100$  MeV
  - These are **Strong Interaction** decays
- Some **much longer lived** hadrons
  - $\tau \sim 10^{-10}$ s
  - Can be measured directly
  - These are **Weak Interaction** decays
- Some with **intermediate lifetimes (e.m.)**



## Conservation Rules

Interaction	Symbol	SI	EM	WI
Energy	E	✓	✓	✓
Momentum	P	✓	✓	✓
Angular Mom <sup>n</sup>	J	✓	✓	✓
Charge (e.m, colour)	Q	✓	✓	✓
Fermion number		✓	✓	✓
Quark number		✓	✓	✓
Baryon number	B	✓	✓	✓
Lepton number	L	✓	✓	✓
Electron number	L <sub>e</sub>	✓	✓	✓
Muon number	L <sub>μ</sub>	✓	✓	✓
Tau number	L <sub>τ</sub>	✓	✓	✓
Quark flavour		✓	✓	✗
Isospin	I	✓	✗	✗
Parity	P	✓	✓	✗
Charge Conjugation	C	✓	✓	✗
Time reversal	T	✓	✓	✗
Matter-Antimatter	CP	✓	✓	✗
Quantum Field Theory	CPT	✓	✓	✓

✓ conserved  
✗ Not necessarily conserved

## [For info.] Running Couplings

EM case

$$\alpha_{EM}(q^2) = \frac{\alpha(0)}{1 - \left(\frac{\alpha(0)}{3\pi}\right) \ln(q^2/m^2)} \quad |q^2| \gg m^2$$

QCD case

$$\alpha_S(q^2) = \frac{\alpha_S(\mu^2)}{1 + \left(\frac{\alpha_S(\mu^2)}{12\pi}\right) [11N_{colours} - 2N_{flavours}] \ln(q^2/m^2)} \quad |q^2| \gg |\mu^2|$$