

Outline

- Relativistic Kinematics
 - ▶ (4-momentum)² invariance, invariant mass
 - ▶ Hypothesis testing, production thresholds
 - ▶ Cross-sections, flux and luminosity, accelerators
 - ▶ Particle lifetime, decay length, width
- Classification of particles
 - ▶ Fermions and bosons
 - ▶ Leptons, hadrons, quarks
 - ▶ Mesons, baryons
- Quark Model
 - ▶ Meson and baryon multiplets
 - ▶ Isospin, strangeness, c, b, t quarks
- Particle Interactions
 - ▶ Virtual particles and range of forces
 - ▶ Strong and weak decays, conservation rules
 - ▶ Parity, charge conjugation, CP
 - ▶ Weak decays of quarks
 - ▶ Colour charge, QCD, gluons
 - ▶ Charmonium and upsilon systems
- Electroweak Interactions
 - ▶ Charged and neutral currents
 - ▶ W, Z, LEP experiments
 - ▶ Higgs and the future
- LHC Experiments
- Future - introduction to accelerator physics

Please see web page for specific references to textbooks and brief reviews from PDG.
(Particularly if you are taking PP Group Studies.)


**Reminder: no lecture on Monday
30 Jan.**
To be re-arranged later in term as required.

Further/Background Reading

Written lecture notes are not distributed but all transparencies/handouts are given below.

- [Lecture 1 \(4 slides/page\)](#) - general introduction to Particle Physics
 - [CERN Summer Student information](#) - deadline 25 Jan. 2012
 - [DESY Summer Student information](#) - deadline 31 Jan. 2012
 - [Review of Quark Model](#), in *2008 Particle Data Group Review of Particle Physics*, C. Amsler et al., published in [Physics Letters B667, 1 \(2008\)](#).
- [Lecture 2 \(4 slides/page\)](#) - Relativistic kinematics and four momenta
 - Griffiths, pages 89-103
 - Williams, page 159
 - [Handout on kinematics and units](#)
 - Units: see also Perkins (4th edition), pg.25.
- [Lecture 3 \(4 slides/page\)](#) - Particle decays and hypothesis testing
 - [Bubble Chamber web \(CERN/G.T.Jones\)](#)
 - [Kinematics. from PDG, K. Nakamura et al. \(Particle Data Group\), J. Phys. G 37, 075021 \(2010\)](#)
- [Lecture 4 \(4 slides/page\)](#) - Fixed target and colliding beam experiments
 - Perkins 4th edition, pp. 28-32 (acceleration in linear vs. circular machines)
 - Perkins 4th edition, pp 32-33 (collider vs. fixed target machines and luminosity)
 - [Table of collider parameters](#):- try to verify luminosity calculation for a few of these?
 - [Brief review of accelerator physics of colliders](#) from Particle Data Group, K. Nakamura et al., JPG 37, 075021 (2010) (<http://pdg.lbl.gov>)
 - See also: Tigner and Chao, [Handbook of Accelerator Physics and Engineering](#) (copy in Library)
 - [USPAS - U.S. Particle Accelerator School](#)
 - [Course material - slides/lecture notes](#)
 - [Joint Accelerator Conference](#) - proceedings for all major accelerator physics conferences

Hypothesis test example



Momenta in GeV/c

	p_x	p_y	p_z
A^+	1.628	0.0897	0
B^-	1.090	-0.0897	0

- Masses m_A, m_B unknown
 ■ 3-momenta inferred from curvature
- Q: How to determine identity of V^0 ?
- We know that
 - ▶ $E_A = (p_A^2 + m_A^2)^{1/2}$ $E_B = (p_B^2 + m_B^2)^{1/2}$
 - ▶ $\underline{p}_V = \underline{p}_A + \underline{p}_B$ $\underline{E}_V = \underline{E}_A + \underline{E}_B$
 - ▶ $m_V = (E_V^2 - p_V^2)^{1/2}$
- Assume A and B are lightest charged hadrons, π^+ , $m_\pi = 139.6 \text{ MeV}/c^2$
- This gives $E_A = (1.628^2 + 0.0897^2 + 0.1396^2)^{1/2} = 1.636 \text{ GeV}$
- $E_B = 1.103 \text{ GeV}$

Hypothesis test example

Momenta in GeV/c

	p_x	p_y	p_z
A^+	1.628	0.0897	0
B^-	1.090	-0.0897	0

- $V^0 \rightarrow A^+ + B^-$
- Assume A and B are lightest charged hadrons, π^+ , $m_\pi = 139.6 \text{ MeV}/c^2$
- This gives $E_A = (1.628^2 + 0.0897^2 + 0.1396^2)^{1/2} = 1.636 \text{ GeV}$
- $E_B = 1.103 \text{ GeV}$
- Using these E_A and E_B , we obtain E_V and \underline{p}_V
 - ▶ $\underline{p}_V = \underline{p}_A + \underline{p}_B$ $E_V = E_A + E_B$
- And from these, it follows that
 - ▶ $m_V = (E_V^2 - \underline{p}_V^2)^{1/2}$
 - ▶ $= ((1.103 + 1.628)^2 - (1.628 + 1.090)^2) = 0.338 \text{ GeV}/c^2$ - unknown!
- Repeat above for A and B = K^+/K^- , $m_K = 494 \text{ MeV}/c^2$
 - ▶ Obtain $m_V = 1.020 \text{ GeV}/c^2$ i.e. $\phi(1020)$ - good hypothesis.

LEP Collider close to max. energy

Beam "lifetime" in e^+e^-

Luminosity vs. time (energy)




Figure 1: Evolution of beam lifetime in LEP.

R. Assmann et al, "Luminosity and Beam Measurements Used for Performance Optimisation in the LEP Collider", EPAC, Vienna, p. 265 (2000).




Figure 2: Luminosity production in 2000. The three angles correspond to 2, 1 and 0 klystrons overhead (right hand numbers, from top to bottom).


G. Arduini et al, "LEP Operation and Performance with 100 GeV Colliding Beams," EPAC, Vienna, p. 265 (2000).

Example: data rates

Physics cross-sections


Integrated collider lumi.

Data corrected for γ radiation



Centre-of-mass energy (GeV)

Integrated lumi. (1/pb)




Days since start up

Example of machine parameters


	CESR (Cornell)	CESR-C (Cornell)	LEP (CERN)	ILC (TBD)
Physics start date	1979	2002	1989	TBD
Physics end date	2002	2008	2000	—
Maximum beam energy (GeV)	6	6	100 - 104.6	250 (upgradeable to 500)
Luminosity ($10^{30} \text{ cm}^{-2}\text{s}^{-1}$)	1280 at 5.3 GeV/beam	76 at 2.08 GeV/beam	24 at Z^0 100 at > 90 GeV	2×10^4
Time between collisions (μs)	0.014 to 0.22	0.014 to 0.22	22	0.3^{\ddagger}
Full crossing angle ($\mu \text{ rad}$)	± 2000	± 3300	0	14000
Energy spread (units 10^{-3})	0.6 at 5.3 GeV/beam	0.82 at 2.08 GeV/beam	0.7→1.5	1
Bunch length (cm)	1.8	1.2	1.0	0.03
Beam radius (μm)	H : 460 V : 4	H : 340 V : 6.5	H : 200 → 300 V : 2.5 → 8	H : 0.639 V : 0.0057
Free space at interaction point (m)	± 2.2 (± 0.6 to REC quads)	± 2.2 (± 0.3 to PM quads)	± 3.5	± 3.5
Luminosity lifetime (hr)	2-3	2-3	20 at Z^0 10 at > 90 GeV	n/a
Turn-around time (min)	5 (topping up)	1.5 (topping up)	50	n/a
Injection energy (GeV)	1.8-6	1.5-6	22	n/a
Transverse emittance ($10^{-9}\pi \text{ rad}\cdot\text{m}$)	H : 210 V : 1	H : 120 V : 3.5	H : 20-45 V : 0.25 → 1	H : 0.02 V : 8×10^{-5} (at 250 GeV)
β^* , amplitude function at interaction point (m)	H : 1.0 V : 0.018	H : 0.94 V : 0.012	H : 1.5 V : 0.05	H : 0.02 V : 0.0004
Beam-beam tune shift per crossing (units 10^{-4})	H : 250 V : 620	e^- : 420 (H), 280 (V) e^+ : 410 (H), 270 (V)	830	n/a
RF frequency (MHz)	500	500	352.2	1300
Particles per bunch (units 10^{10})	1.15	4.7	45 in collision 60 in single beam	2
Bunches per ring per species	9 trains of 5 bunches	8 trains of 3 bunches	4 trains of 1 or 2	2625

Higher energy e^+e^- colliders

- In planning/R&D phase
- Physics motivations many
- Cross-sections small!
- Luminosity **the** issue!
- b/c-tagging with high purity/efficiency
 - ▶ e.g. Higgs branching ratios
- Precision Tracking
- Recoil mass measurements
- Jet energy resolution
- Multi jet final states, e.g.
 - ▶ $t\bar{t}$
 - ▶ separation of WW/ZZ



Beam-Beam effects in future e^+e^- machines



- Mutual focussing, “pinch” enhancement
- Large disruption – single pass machine
- Beam-beam effects, amplify initial vertical offsets

▶ detect downstream, feedback, maintains lumi

Luminosity in future e^+e^- machine

■ High luminosity achieved by

[c/o Andrei Seryi]

- ▶ Many incident particles
- ▶ Small transverse cross-section at interaction point



■ e.g., LC beam sizes just before collision (500 GeV):

250 * 3 * 110000 nm

(x y z)



vertical size
is smallest



$$\frac{\Delta E}{E} \propto \frac{E_{CM}}{\sigma_z} \frac{N^2}{(\sigma_x + \sigma_y)^2}$$

$$L = \frac{f_{rep} n_b N^2}{4\pi \sigma_x \sigma_y} H_D$$