

Outline

- Relativistic Kinematics
 - ▶ (4-momentum)² invariance, invariant mass
 - ▶ Hypothesis testing, production thresholds
 - ▶ Cross-sections, flux and luminosity, accelerators
 - ▶ Particle lifetime, decay length, width
- Classification of particles
 - ▶ Fermions and bosons
 - ▶ Leptons, hadrons, quarks
 - ▶ Mesons, baryons
- Quark Model
 - ▶ Meson and baryon structure
 - ▶ Isospin, strangeness
- Particle Interactions
 - ▶ Virtual particles
 - ▶ Strong and weak interactions
 - ▶ Parity, charge conjugation
 - ▶ Weak decays
 - ▶ Colour charge, QCD, gluons
 - ▶ Charmonium and upsilon systems
- Electroweak Interactions
 - ▶ Charged and neutral currents
 - ▶ W, Z, LEP
 - ▶ Higgs and top quark
- LHC Experiments
- Future - introduction to accelerator physics

Today

- Lecture 4 (4 slides/page) - Fixed target and colliding beam experiments
 - Perkins 3rd edition, pp. 28-32 (acceleration in linear vs. circular machines)
 - Perkins 3rd edition, pp. 32-33 (collider vs. fixed target machines and luminosity)
 - Table of collider parameters - try to verify luminosity calculation for a few of these?
 - Brief review of accelerator physics of colliders from Particle Data Group, K. Nakamura et al., JPG
 - See also: Tigner and Chao, Handbook of Accelerator Physics and Engineering (copy in Library)
 - LSPAS - U.S. Particle Accelerator School
 - Course material - slides/lecture notes
 - Joint Accelerator Conference - proceedings for all major accelerator physics conferences

Previous Lecture

- Lecture 3 (4 slides/page) - Particle decays and hypothesis testing
 - Bubble Chamber web (CERN/IG T.Jones)
 - Kinematics from PDG, J. Berninger et al. (Particle Data Group), Phys. Rev. D88, 010001 (2012)

Further/Background Reading

Y3 Particle Physics

Material for the 2012-13 course:

- Lectures start 7 January 2013 and are:
 - Mondays at 1500, Walton Lecture Theatre C
 - Thursdays 1500, Physics Process Small Lecture Theatre (C00)

References for further reading/background information will be given below each lecture.

The course [Study List](#) is similar to previous PP courses.

- Course Summary
- Study List
- Useful additional collection of web pages

Written lecture notes are not distributed but all transparencies/handouts are given below:

- Lecture 1 (4 slides/page) - general introduction to Particle Physics
- [Review of Quark Models](#) - in 2009 Particle Data Group Review of Particle Physics, C. Amsler et al., published in Physics Letters B067, 1 (2008)
- CERN Summer Student Lectures
 - CERN Summer Student information - deadline 27 Jan. 2013
 - CERN Summer student passes - [return here to see](#)
 - CERN Summer Student info
 - CERN Summer Student information - application deadline 31 Jan. 2013
- Lecture 2 (4 slides/page) - Relativistic kinematics and four momenta
 - Griffiths, page 89-103
 - Williams, page 159
 - Handout on relativistic kinematics
 - Links: see also Perkins (3rd edition), pg 26
- Lecture 3 (4 slides/page) - Particle decay and hypothesis testing

Per lecture reading in course texts – please look at it
Library reading list for course updated/expanded

LEP Collider close to max. energy

Beam "lifetime" in e^+e^- Luminosity vs. time (energy)




Figure 1: Evolution of beam lifetime in LEP.

R. Assmann et al, "Luminosity and Beam Measurements Used for Performance Optimisation in the LEP Collider", EPAC, Vienna, p. 265 (2000).






Figure 2: Luminosity production in 2000. The three angles correspond to 2, 1 and 0 klystrons overhead (right hand numbers, from top to bottom).

G. Arduini et al, "LEP Operation and Performance with 100 GeV Colliding Beams," EPAC, Vienna, p. 265 (2000).

Example: data rates

Physics cross-sections Integrated collider lumi.



Days since start up


Centre-of-mass energy (GeV)

Example of machine parameters


	CERN (CERN)	CERN (CERN)	LEP (CERN)	ILC (FBI)
Physics start date	1979	2002	1990	TBD
Physics end date	2002	2008	2000	—
Maximum beam energy (GeV)	0	6	100–134.6	250 (approachable to 500)
Luminosity (10^{34} cm $^{-2}$ s $^{-1}$)	1200 at 5.3 GeV/beam	70 at 2.08 GeV/beam	24 at 2^{nd} 100 at > 50 GeV	2×10^6
Time between collisions (ps)	0.014 to 0.22	0.014 to 0.22	22	0.37
Full crossing angle (μrad)	±2000	±2300	0	14000
Energy spread (units 10^{-4})	0.6 at 5.3 GeV/beam	0.62 at 2.08 GeV/beam	0.7–1.5	1
Bunch length (mm)	1.8	1.2	1.0	0.03
Beam radius (μm)	$R: 600$ $V: 4$	$R: 140$ $V: 6.5$	$R: 200–300$ $V: 2.5–8$	$R: 0.620$ $V: 0.0057$
Free space at interaction point (m)	3.27 (±0.6, no PMA quadr.)	3.27 (±0.3, no PMA quadr.)	±3.5	±3.5
Luminosity lifetime (hr)	2–3	2–3	20 at 2^{nd} 10 at > 50 GeV	n/a
Transverse time (ns)	5 (trapping up)	1.5 (trapping up)	30	n/a
Injection energy (GeV)	1.8–6	1.5–6	22	n/a
Transverse emittance (10^{-9} e.m.u.)	$R: 210$ $V: 1$	$R: 120$ $V: 3.5$	$R: 20–45$ $V: 0.25–1$	$R: 0.02$ $V: 4 \times 10^{-5}$ (at 250 GeV)
β^* , magnitude function at interaction point (m)	$R: 1.0$ $V: 0.018$	$R: 0.94$ $V: 0.012$	$R: 1.5$ $V: 0.05$	$R: 0.02$ $V: 0.0004$
Beam-beam tune shift per crossing (units 10^{-4})	$R: 250$ $V: 520$	~ 520 ($R: 240$ V)	800	n/a
RF frequency (MHz)	500	500	302.2	1300
Particles per bunch (units 10^{11})	1.15	4.7	43 in collision 40 in single beam	2
Bunches per ring per species	9 trunks of 5 bunches	8 trunks of 3 bunches	4 trunks of 1 or 2	2025

Higher energy e^+e^- colliders

- In planning/R&D phase
- Physics motivations many
- Cross-sections small!
- Luminosity the issue!
- b/c-tagging with high purity/efficiency
 - ▶ e.g. Higgs branching ratios
- Precision Tracking
- Recoil mass measurements
- Jet energy resolution
- Multi jet final states, e.g.
 - ▶ $t\bar{t}$ bar
 - ▶ separation of WW/ZZ



Beam-Beam effects in future e^+e^- machines



- Mutual focussing, “pinch” enhancement
- Large disruption – single pass machine
- Beam-beam effects, amplify initial vertical offsets
 - ▶ detect downstream, feedback, maintains lumi

Glasgow, 24-Jan-2008


Nigel Watson / Birmingham

Luminosity in future e^+e^- machine

- High luminosity achieved by
 - ▶ Many incident particles
 - ▶ Small transverse cross-section at interaction point

e.g., LC beam sizes just before collision (500 GeV):
 $250 \times 3 \times 110000$ nm

(x y z)
 vertical size is smallest



$$\frac{\Delta E}{E} \propto \frac{E_{CM}}{\sigma_z} \frac{N^2}{(\sigma_x + \sigma_y)^2}$$

$$L = \frac{f_{rep} n_b N^2}{4\pi \sigma_x \sigma_y} H_D$$

Glasgow, 24-Jan-2008

Nigel Watson / Birmingham